



(Callum Weir)

## Reduced tillage - Non-inversion and shallow cultivation in organic systems (United Kingdom)

Non-inversion and shallow cultivation in organic systems

### DESCRIPTION

**Non-inversion and 'shallow' ploughing cultivation strategies on an organic farm, where the use of herbicides for weed control is prohibited.**

The shallow plough is used on land dominated by clay soils at an organically farmed estate, south of Cambridgeshire, UK. Previously, 'conventional' ploughs were used, which plough deeper than a shallow plough. However, ploughing deeper would often bring large chunks of raw clay from the subsoil to the surface. This would quickly solidify, locally referred to as when the soil turns to 'concrete'. Numerous cultivations were then required to reduce these 'concrete' soil chunks into a seed bed. It was a laborious, expensive task which sacrificed soil health to produce a less than satisfactory result. However, the farm still required a plough of some form as a means of weed control through inversion. As it is an organic estate, chemical sprays could not be used. A shallow plough was invested in as a way of striking the balance between overcoming the problems of creating a seedbed, but also maintaining the weed control benefits of inversion tillage. It has been very successful in reducing the input requirements, and at the same time increasing the quality of the output. Whilst shallow ploughing has challenges, such as full inversion of weeds in very dry conditions, on balance it is much better for the farming business than the previous alternative. We are able to do less damage to soil, and increase outputs which is important due to agricultural labour scarcity and smaller weather windows due to climate change.

Reduced tillage options have been a challenge to combat in organic systems where herbicides are prohibited. As such, trials of reduced tillage options have been explored. These include;

- 1) Non-inversion tillage where no ploughing is done and soil is cultivated to the first 100 mm.
- 2) Shallow ploughing where a specifically designed plough inverts soil to a depth of 125 mm, as opposed to traditional plough depths of 200 mm.


The purpose of this technology is to minimise soil disturbance to enhance the soil structure, biology and chemistry, whilst creating a seed bed and controlling weeds. The challenge on the specific site is there has been a history of annual plough, which has led to the proliferation of weeds that thrive on such systems. These include creeping thistle and common docks. As such, there was also the purpose of 'disrupting' the existing system in order to control these weeds. The only specific input required was a shallow plough, designed to invert soil from lower depths. For non-inversion tillage, a subsoiler and disc cultivator were used. The non-inversion tillage was done at two sites; one cereal stubble and one out of a fertility building two-year grass and clover ley.

Benefits/impacts/things land owners did/did not like:

Non-inversion tillage:

- Instead of ploughing, non-inversion tillage from the fertility ley allowed us to keep the soil structure from 2 years of grass/clover intact and in the right soil profile. We weren't burying the friable, high-nutrient and porous top soil 200 mm under the ground and we weren't lifting heavy, lower-aerobic soil to the surface where we wanted to plant.
- This meant that plants established quicker and we were able to drill later, despite the

### LOCATION



**Location:** Wimpole Estate, United Kingdom

**No. of Technology sites analysed:** single site

**Geo-reference of selected sites**

- -0.04205, 52.14849
- -0.04205, 52.14849

**Spread of the Technology:** evenly spread over an area (4.0 km<sup>2</sup>)

**In a permanently protected area?:** No

**Date of implementation:** 2018; less than 10 years ago (recently)

**Type of introduction**

- through land users' innovation
- as part of a traditional system (> 50 years)
- during experiments/ research
- through projects/ external interventions

fields being very heavy, poorly drained fields.

-Weeds were killed, primarily through timely cultivations during a hot-spell, so that the cultivator brought roots to the surface to dry them.

-Drainage was evident after drilling as we were able to graze sheep on the wheat in March.

-Crops have tillered well and responded to nutrients.

-Establishment costs were approximately £30/ha cheaper.

-However, non-inversion tillage in cereal stubbles has not been as successful due to weed control, and whether the cheaper costs outweighs the weed burden remains to be assessed. The reason for this is not being able to cultivate during the hot weather (as this came before harvest).

-In addition, in cereal stubbles, we have seen less creeping thistles and docks, but more wild oats and cereal volunteers.

Shallow ploughing:

-Cheaper establishment costs through lower diesel usage (yet to be quantified).

-Better in many circumstances of inverting soil completely, but from a much lower depth.

-Did not bring up any large clumps of sub-soil which the conventional plough would.

These result in much cultivations to break the clumps down.

-Ploughing 'on-land' meant that there was no smearing in the furrow from tyres.

-Lower HP requirement – 180 hp tractor ploughing 3.2 m to 125 mm on heavy land.

-Ploughing left over-winter did not require more than one cultivation before drilling as ploughed soil was friable from lower plough depth.

-That being said, there were favourable ploughing conditions in 2018. Regardless, we have sold our conventional plough because we like the shallow plough so much.

General benefits are:

-Reduced, prevented or restored land degradation

-Improved/preserved biodiversity

-Increased adaptation/resilience to climate change/extremes and its impacts

-A potential beneficial economic impact

The compilation of this SLM is a part of the European Interreg project FABulous Farmers which aims to reduce the reliance on external inputs by encouraging the use of methods and interventions that increase the farm's Functional AgroBiodiversity (FAB). Visit [www.fabulousfarmers.eu](http://www.fabulousfarmers.eu) and [www.nweurope.eu/Fabulous-Farmers](http://www.nweurope.eu/Fabulous-Farmers) for more information.



Shallow ploughing green stubbles. (Callum Weir)



Claas Arion tractor ploughing (Callum Weir)

## CLASSIFICATION OF THE TECHNOLOGY

### Main purpose

- improve production
- reduce, prevent, restore land degradation
- conserve ecosystem
- protect a watershed/ downstream areas – in combination with other Technologies
- preserve/ improve biodiversity
- reduce risk of disasters
- adapt to climate change/ extremes and its impacts
- mitigate climate change and its impacts
- create beneficial economic impact
- create beneficial social impact

### Land use

Land use mixed within the same land unit: No



### Cropland

- Annual cropping: cereals - other, fodder crops - clover, fodder crops - grasses
- Number of growing seasons per year: 1  
Is intercropping practiced? Yes  
Is crop rotation practiced? No

### Water supply

- rainfed
- mixed rainfed-irrigated
- full irrigation

### Purpose related to land degradation

- prevent land degradation
- reduce land degradation
- restore/ rehabilitate severely degraded land
- adapt to land degradation
- not applicable

### Degradation addressed



**soil erosion by wind** - Et: loss of topsoil



**chemical soil deterioration** - Cn: fertility decline and reduced organic matter content (not caused by erosion)



**physical soil deterioration** - Pc: compaction



**biological degradation** - Bs: quality and species composition/ diversity decline, Bp: increase of pests/ diseases, loss of predators



**water degradation** - Hs: change in quantity of surface water, Hg: change in groundwater/aquifer level, Hq: decline of groundwater quality, Hw: reduction of the buffering capacity of wetland areas

### SLM group

- minimal soil disturbance
- integrated soil fertility management
- water diversion and drainage

### SLM measures



**agronomic measures** - A3: Soil surface treatment (A 3.2: Reduced tillage (> 30% soil cover)), A5: Seed management, improved varieties

## TECHNICAL DRAWING

### Technical specifications

## ESTABLISHMENT AND MAINTENANCE: ACTIVITIES, INPUTS AND COSTS

### Calculation of inputs and costs

- Costs are calculated: per Technology area (size and area unit: 4 ha; conversion factor to one hectare: 1 ha = Approx. £45/ha – about 25% less than 'deep ploughing')
- Currency used for cost calculation: **GBP**
- Exchange rate (to USD): 1 USD = 0.82 GBP
- Average wage cost of hired labour per day: £90

### Most important factors affecting the costs

Most important factors affecting cost are decreased time spent ploughing and lower diesel cost, reducing establishment costs by £15 per ha.

### Establishment activities

1. Use of shallow plough (Timing/ frequency: After harvest)

### Establishment inputs and costs (per 4 ha)

Specify input	Unit	Quantity	Costs per Unit (GBP)	Total costs per input (GBP)	% of costs borne by land users
<b>Labour</b>					
Person per day	person day	1.0	90.0	90.0	100.0
<b>Equipment</b>					
Ovlac Shallow Plough (7+1f) (one off)	1	1.0	11000.0	11000.0	100.0
Tractor	per day	1.0	180.0	180.0	100.0
<b>Other</b>					
Diesel (120 litres per day)	ltrs per day	1.0	60.0	60.0	100.0
<b>Total costs for establishment of the Technology</b>				<b>11'330.0</b>	
<i>Total costs for establishment of the Technology in USD</i>				<i>13'817.07</i>	

### Maintenance activities

1. Grease plough (Timing/ frequency: once per week)
2. change plough points (Timing/ frequency: once per season)

## NATURAL ENVIRONMENT

### Average annual rainfall

- < 250 mm
- 251-500 mm
- 501-750 mm
- 751-1,000 mm
- 1,001-1,500 mm
- 1,501-2,000 mm
- 2,001-3,000 mm
- 3,001-4,000 mm
- > 4,000 mm

### Agro-climatic zone

- humid
- sub-humid
- semi-arid
- arid

### Specifications on climate

Highest rainfall month is August, which is important as this is when cultivations need to occur. As non-inversion and shallow ploughing are faster operations, this means that cultivations can occur at more optimum times.

### Slope

- flat (0-2%)
- gentle (3-5%)
- moderate (6-10%)
- rolling (11-15%)

### Landforms

- plateau/plains
- ridges
- mountain slopes
- hill slopes

### Altitude

- 0-100 m a.s.l.
- 101-500 m a.s.l.
- 501-1,000 m a.s.l.
- 1,001-1,500 m a.s.l.

### Technology is applied in

- convex situations
- concave situations
- not relevant

<input type="checkbox"/> hilly (16-30%) <input type="checkbox"/> steep (31-60%) <input type="checkbox"/> very steep (>60%)	<input type="checkbox"/> footslopes <input type="checkbox"/> valley floors	<input type="checkbox"/> 1,501-2,000 m a.s.l. <input type="checkbox"/> 2,001-2,500 m a.s.l. <input type="checkbox"/> 2,501-3,000 m a.s.l. <input type="checkbox"/> 3,001-4,000 m a.s.l. <input type="checkbox"/> > 4,000 m a.s.l.	
<b>Soil depth</b> <input type="checkbox"/> very shallow (0-20 cm) <input type="checkbox"/> shallow (21-50 cm) <input checked="" type="checkbox"/> moderately deep (51-80 cm) <input type="checkbox"/> deep (81-120 cm) <input type="checkbox"/> very deep (> 120 cm)	<b>Soil texture (topsoil)</b> <input type="checkbox"/> coarse/ light (sandy) <input type="checkbox"/> medium (loamy, silty) <input checked="" type="checkbox"/> fine/ heavy (clay)	<b>Soil texture (&gt; 20 cm below surface)</b> <input type="checkbox"/> coarse/ light (sandy) <input type="checkbox"/> medium (loamy, silty) <input checked="" type="checkbox"/> fine/ heavy (clay)	<b>Topsoil organic matter content</b> <input checked="" type="checkbox"/> high (>3%) <input type="checkbox"/> medium (1-3%) <input type="checkbox"/> low (<1%)
<b>Groundwater table</b> <input type="checkbox"/> on surface <input checked="" type="checkbox"/> < 5 m <input type="checkbox"/> 5-50 m <input type="checkbox"/> > 50 m	<b>Availability of surface water</b> <input type="checkbox"/> excess <input type="checkbox"/> good <input checked="" type="checkbox"/> medium <input type="checkbox"/> poor/ none	<b>Water quality (untreated)</b> <input type="checkbox"/> good drinking water <input type="checkbox"/> poor drinking water (treatment required) <input type="checkbox"/> for agricultural use only (irrigation) <input type="checkbox"/> unusable <i>Water quality refers to: both ground and surface water</i>	<b>Is salinity a problem?</b> <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No  <b>Occurrence of flooding</b> <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No
<b>Species diversity</b> <input checked="" type="checkbox"/> high <input type="checkbox"/> medium <input type="checkbox"/> low	<b>Habitat diversity</b> <input checked="" type="checkbox"/> high <input type="checkbox"/> medium <input type="checkbox"/> low		

### CHARACTERISTICS OF LAND USERS APPLYING THE TECHNOLOGY

<b>Market orientation</b> <input type="checkbox"/> subsistence (self-supply) <input type="checkbox"/> mixed (subsistence/ commercial) <input checked="" type="checkbox"/> commercial/ market	<b>Off-farm income</b> <input checked="" type="checkbox"/> less than 10% of all income <input type="checkbox"/> 10-50% of all income <input type="checkbox"/> > 50% of all income	<b>Relative level of wealth</b> <input type="checkbox"/> very poor <input type="checkbox"/> poor <input checked="" type="checkbox"/> average <input type="checkbox"/> rich <input type="checkbox"/> very rich	<b>Level of mechanization</b> <input type="checkbox"/> manual work <input type="checkbox"/> animal traction <input checked="" type="checkbox"/> mechanized/ motorized
<b>Sedentary or nomadic</b> <input checked="" type="checkbox"/> Sedentary <input type="checkbox"/> Semi-nomadic <input type="checkbox"/> Nomadic	<b>Individuals or groups</b> <input checked="" type="checkbox"/> individual/ household <input type="checkbox"/> groups/ community <input type="checkbox"/> cooperative <input type="checkbox"/> employee (company, government)	<b>Gender</b> <input type="checkbox"/> women <input checked="" type="checkbox"/> men	<b>Age</b> <input type="checkbox"/> children <input type="checkbox"/> youth <input checked="" type="checkbox"/> middle-aged <input checked="" type="checkbox"/> elderly
<b>Area used per household</b> <input type="checkbox"/> < 0.5 ha <input type="checkbox"/> 0.5-1 ha <input type="checkbox"/> 1-2 ha <input type="checkbox"/> 2-5 ha <input type="checkbox"/> 5-15 ha <input type="checkbox"/> 15-50 ha <input type="checkbox"/> 50-100 ha <input checked="" type="checkbox"/> 100-500 ha <input checked="" type="checkbox"/> 500-1,000 ha <input type="checkbox"/> 1,000-10,000 ha <input type="checkbox"/> > 10,000 ha	<b>Scale</b> <input type="checkbox"/> small-scale <input checked="" type="checkbox"/> medium-scale <input type="checkbox"/> large-scale	<b>Land ownership</b> <input type="checkbox"/> state <input type="checkbox"/> company <input type="checkbox"/> communal/ village <input type="checkbox"/> group <input type="checkbox"/> individual, not titled <input checked="" type="checkbox"/> individual, titled	<b>Land use rights</b> <input type="checkbox"/> open access (unorganized) <input type="checkbox"/> communal (organized) <input checked="" type="checkbox"/> leased <input checked="" type="checkbox"/> individual  <b>Water use rights</b> <input type="checkbox"/> open access (unorganized) <input type="checkbox"/> communal (organized) <input type="checkbox"/> leased <input type="checkbox"/> individual

### Access to services and infrastructure

health	poor	<input checked="" type="checkbox"/>	good
education	poor	<input checked="" type="checkbox"/>	good
technical assistance	poor	<input checked="" type="checkbox"/>	good
employment (e.g. off-farm)	poor	<input checked="" type="checkbox"/>	good
markets	poor	<input checked="" type="checkbox"/>	good
energy	poor	<input checked="" type="checkbox"/>	good
roads and transport	poor	<input checked="" type="checkbox"/>	good
drinking water and sanitation	poor	<input checked="" type="checkbox"/>	good
financial services	poor	<input checked="" type="checkbox"/>	good

### IMPACTS

#### Socio-economic impacts

Crop production	decreased	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	increased
crop quality	decreased	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	increased
fodder production	decreased	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	increased
animal production	decreased	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	increased

Crop quality before SLM not able to handle grazing, but now can graze so large increase in fodder/animal production compared to previous model

Crop quality before SLM not able to handle grazing,

but now can graze so large increase in fodder/animal production compared to previous model

land management	hindered		✓		simplified
expenses on agricultural inputs	increased		✓		decreased
farm income	decreased		✓		increased
diversity of income sources	decreased		✓		increased
workload	increased		✓		decreased
work/life balance	None		✓		None

**Socio-cultural impacts**

recreational opportunities	reduced		✓		improved
SLM/ land degradation knowledge	reduced		✓		improved

**Ecological impacts**

harvesting/ collection of water (runoff, dew, snow, etc)	reduced		✓		improved
surface runoff	increased		✓		decreased
excess water drainage	reduced		✓		improved
soil moisture	decreased		✓		increased
soil cover	reduced		✓		improved
soil loss	increased		✓		decreased
soil crusting/ sealing	increased		✓		reduced
soil compaction	increased		✓		reduced
nutrient cycling/ recharge	decreased		✓		increased
soil organic matter/ below ground C	decreased		✓		increased
vegetation cover	decreased		✓		increased
biomass/ above ground C	decreased		✓		increased
invasive alien species	increased		✓		reduced
animal diversity	decreased		✓		increased
beneficial species (predators, earthworms, pollinators)	decreased		✓		increased
pest/ disease control	decreased		✓		increased

A small decrease in disease control with shallow ploughing is not as effective as inverting with a conventional plough. This is because less of the stubble from the previous crop would be inverted, creating a greater chance of disease carryover, for example Septoria nodorum blotch.

flood impacts	increased		✓		decreased
drought impacts	increased		✓		decreased

**Off-site impacts**

reliable and stable stream flows in dry season (incl. low flows)	reduced		✓		increased
downstream flooding (undesired)	increased		✓		reduced
downstream siltation	increased		✓		decreased
groundwater/ river pollution	increased		✓		reduced
buffering/ filtering capacity (by soil, vegetation, wetlands)	reduced		✓		improved
wind transported sediments	increased		✓		reduced
impact of greenhouse gases	increased		✓		reduced

### COST-BENEFIT ANALYSIS

**Benefits compared with establishment costs**

Short-term returns	very negative		✓		very positive
Long-term returns	very negative		✓		very positive

**Benefits compared with maintenance costs**

Short-term returns	very negative		✓		very positive
Long-term returns	very negative		✓		very positive

### CLIMATE CHANGE










**Gradual climate change**

annual temperature increase	not well at all		✓		very well	
seasonal temperature increase	not well at all		✓		very well	Season: spring
annual rainfall decrease	not well at all		✓		very well	
seasonal rainfall decrease	not well at all		✓		very well	Season: spring

**Climate-related extremes (disasters)**







local rainstorm	not well at all		✓		very well	
-----------------	-----------------	--	---	--	-----------	--

local thunderstorm  
heatwave  
drought

not well at all     very well  
not well at all     very well  
not well at all     very well

#### Other climate-related consequences

extended growing period  
reduced growing period

not well at all     very well  
not well at all     very well

## ADOPTION AND ADAPTATION

### Percentage of land users in the area who have adopted the Technology

- single cases/ experimental
- 1-10%
- 11-50%
- > 50%

### Of all those who have adopted the Technology, how many have done so without receiving material incentives?

- 0-10%
- 11-50%
- 51-90%
- 91-100%

### Has the Technology been modified recently to adapt to changing conditions?

- Yes
- No

### To which changing conditions?

- climatic change/ extremes
- changing markets
- labour availability (e.g. due to migration)

## CONCLUSIONS AND LESSONS LEARNT

### Strengths: land user's view

- Cheaper establishment costs and quicker establishment time mean it will benefit the farm in the long term as labour becomes an issue (regardless of Brexit).
- Makes soil more resilient to changing weather conditions, both drier and wetter conditions.
- Reduced soil carbon emissions and diesel emissions from tractor.
- Better soil structure, biology and chemistry to boost yield, plus allows us to use plough sparingly as a 'reset' button when we really need to.
- However, there is a risk to yield if not used correctly. Plus, we may solve one weed issue (thistles and docks) and move to another weed issue (cereal volunteers, blackgrass and wild oats).

### Strengths: compiler's or other key resource person's view

### Weaknesses/ disadvantages/ risks: land user's view → how to overcome

- We may solve one weed issue (thistles and docks) and move to another weed issue (cereal volunteers, blackgrass and wild oats). → - Use dry June/July to non-invert fertility leys, allowing plough to be used as a reset button later in the rotation.
  - Minimise non-inversion in cereal stubbles to cleanest crops.
- Management demand to adapt technology to annual changes in conditions (not as easy as ploughing or spraying in any conditions - to do this, you must be adaptable). → -Operator education
  - Planning

### Weaknesses/ disadvantages/ risks: compiler's or other key resource person's view → how to overcome

## REFERENCES

### Compiler

Sabine Reinsch

Date of documentation: June 10, 2019

### Resource persons

Callum Weir - land user

### Full description in the WOCAT database

[https://qcat.wocat.net/en/wocat/technologies/view/technologies\\_5012/](https://qcat.wocat.net/en/wocat/technologies/view/technologies_5012/)

### Linked SLM data

n.a.

### Documentation was facilitated by

Institution

- The National Trust (National Trust) - United Kingdom
- UK Centre for Ecology & Hydrology (CEH) - United Kingdom

Project

- European Interreg project FABulous Farmers

### Reviewer

Renate Fleiner

Last update: Nov. 29, 2019